



Results from the national surveillance of SARS-CoV-2 in wastewater

Content

Information about the national wastewater surveillance of SARS-CoV-2	2
Where do we measure wastewater concentrations of SARS-CoV-2?	2
How do we measure wastewater concentrations of SARS-CoV-2?	2
How are the results of wastewater measurements presented?	2
 Surveillance of wastewater concentrations of SARS-CoV-2, week 42	 3
Denmark	3
Capital Region of Denmark	7
Central Denmark Region	10
North Denmark Region	14
Region Zealand	17
Region of Southern Denmark	20



Information about the national wastewater surveillance of SARS-CoV-2

Below is a short description of the wastewater surveillance of SARS-CoV-2 in Denmark. A detailed description of the wastewater surveillance can be found on SSI's website ([link to detailed description](#)).

Where do we measure wastewater concentrations of SARS-CoV-2?

In the national wastewater surveillance of SARS-CoV-2, 29 wastewater samples are taken from 28 treatment plants across Denmark.

How do we measure wastewater concentrations of SARS-CoV-2?

Genetic material (RNA) of the SARS-CoV-2 virus are excreted in the feces of approximately half of the infected individuals and can therefore be measured in wastewater. Wastewater samples are transported to SSI's laboratory, where they are preprocessed and analyzed using PCR tests (RT-qPCR). This provides an estimate of the number of RNA copies of SARS-CoV-2 per liter of wastewater. The PCR analysis also includes the naturally occurring virus PMMoV, that is excreted in the feces. The laboratory results are analyzed by the Infectious Disease Epidemiology & Prevention department at SSI.

How are the results of wastewater measurements presented?

There are two main categories of wastewater results: A weekly weighted average of the SARS-CoV-2 concentration in wastewater, and a growth rate that describes the change in the national concentration (increasing, stable, or decreasing) based on the three most recent weeks of wastewater data.


The weekly weighted average for the last 15 months is presented nationally and for each region. For each sampling site, the weekly weighted average is shown after April 1, 2023. The results are presented first for Denmark and then for each region in alphabetical order.

Surveillance of wastewater concentrations of SARS-CoV-2, week 42

Denmark

The figure below shows the concentration of SARS-CoV-2 in the wastewater, aggregated for all the sampling sites in Denmark. The figure also shows the national number of new hospital admissions due to SARS-CoV-2.

Wastewater concentration of SARS-CoV-2






The figure below shows a classification of the SARS-CoV-2 concentration in the wastewater samples. There are two thresholds: LoD (Limits of Detection) and LoQ (Limit of Quantification). Samples classified as < LoD have a concentration of SARS-CoV-2 that is lower than the detection limit of the method. Samples classified as < LoQ have a concentration of SARS-CoV-2 that is high enough to be detected with 95 % certainty, but too low to be quantified. The quantified samples are those that have a concentration of SARS-CoV-2 that is high enough to quantify and the missing samples are those that for various reasons are not included in the analysis.

In more than two of the past three weeks, more than 2/3 of the wastewater samples were classified as "Quantified". Hence, the concentration of SARS-CoV-2 in the wastewater is at a level that is high enough for growth rates to be calculated and published.


Categorization of quantifiability



The average weekly growth rate in Denmark, based on observations from the past three weeks, is shown below.


Week 42: Tendency in wastewater surveillance

Percentwise change based on the measurements from the last three weeks





A map of the catchment areas of the included wastewater treatment plants is seen below.






Capital Region of Denmark


The wastewater concentration of SARS-CoV-2 in the Capital Region of Denmark, aggregated and for each sampling site, is shown below.

Wastewater concentration of SARS-CoV-2, Capital Region







Wastewater concentration of SARS-CoV-2, Hillerød (R)



Wastewater concentration of SARS-CoV-2, Hvidovre (Avedøre) (R)




Wastewater concentration of SARS-CoV-2, København (Damhusåen) (R)





Wastewater concentration of SARS-CoV-2, Lynetten (nordre tilløb) (R)




Number of inhabitants: 213.193

Wastewater concentration of SARS-CoV-2, Lynetten (søndre tilløb) (R)



Number of inhabitants: 441.519

Wastewater concentration of SARS-CoV-2, Måløv (R)




Number of inhabitants: 51.329

Central Denmark Region


The wastewater concentration of SARS-CoV-2 in the Central Denmark Region, aggregated and for each sampling site, is shown below.

Wastewater concentration of SARS-CoV-2, Central Denmark Region







Wastewater concentration of SARS-CoV-2, Aarhus (Egå) (R)




Wastewater concentration of SARS-CoV-2, Aarhus (Marselisborg) (R)



Wastewater concentration of SARS-CoV-2, Aarhus (Viby) (R)




Wastewater concentration of SARS-CoV-2, Aarhus (Åby) (R)




Number of inhabitants: 60.662

Wastewater concentration of SARS-CoV-2, Herning (R)



Number of inhabitants: 56.073

Wastewater concentration of SARS-CoV-2, Horsens (R)




Number of inhabitants: 81.018

25.10.2023




Wastewater concentration of SARS-CoV-2, Søholt (R)



North Denmark Region


The wastewater concentration of SARS-CoV-2 in the North Denmark Region, aggregated and for each sampling site, is shown below.

Wastewater concentration of SARS-CoV-2, North Denmark Region







Wastewater concentration of SARS-CoV-2, Aalborg Vest (R)



Wastewater concentration of SARS-CoV-2, Aalborg Øst (R)




Wastewater concentration of SARS-CoV-2, Hjørring (R)





Wastewater concentration of SARS-CoV-2, Mariagerfjord (R)







Region Zealand

The wastewater concentration of SARS-CoV-2 in Region Zealand, aggregated and for each sampling site, is shown below.

Wastewater concentration of SARS-CoV-2, Region Zealand




Wastewater concentration of SARS-CoV-2, Holbæk (R)




Number of inhabitants: 36.955

Wastewater concentration of SARS-CoV-2, Køge (R)



Number of inhabitants: 54.139


Wastewater concentration of SARS-CoV-2, Mosehede (R)




Number of inhabitants: 48.406




Wastewater concentration of SARS-CoV-2, Næstved (R)



Wastewater concentration of SARS-CoV-2, Ringsted C (R)



Wastewater concentration of SARS-CoV-2, Slagelse (R)







Region of Southern Denmark

The wastewater concentration of SARS-CoV-2 in the Region of Southern Denmark, aggregated and for each sampling site, is shown below.

Wastewater concentration of SARS-CoV-2, Region of Southern Denmark




Wastewater concentration of SARS-CoV-2, Esbjerg Vest (R)




Number of inhabitants: 62.097

Wastewater concentration of SARS-CoV-2, Fredericia (R)




Number of inhabitants: 52.187

Wastewater concentration of SARS-CoV-2, Kolding (R)




Number of inhabitants: 73.687

Wastewater concentration of SARS-CoV-2, Odense (Ejby Mølle) (R)



Wastewater concentration of SARS-CoV-2, Odense (Nordvest) (R)



Wastewater concentration of SARS-CoV-2, Vejle (R)

